#### **IUSTI**

# 2013 European Guideline on the Management of Lymphogranuloma Venereum

Date: 30/01/2013

Henry JC de Vries<sup>1</sup>, Adele Zingoni<sup>2</sup>, Alexander Kreuter<sup>3</sup>, Harald Moi<sup>4</sup>, John A White<sup>5</sup>
1 STI outpatient clinic, Cluster Infectious Diseases, Health Service Amsterdam, Amsterdam, The Netherlands; Department of Dermatology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands; Centre for Infectious Diseases and Immunology Amsterdam (CINIMA), Amsterdam, The Netherlands; Centre for Infectious Disease Control, National Institute of Public Health and the Environment, Bilthoven, the Netherlands

- 2 Department of Biomedical Sciences and Human Oncology, Dermatologic Clinic, University of Turin, Turin, Italy
- 3 Department of Dermatology, Venereology, and Allergology, HELIOS St. Elisabeth Hospital Oberhausen, Germany
- 4 Olafia Clinic, Oslo university Hospital, Institute of Clinical Medicine, University of Oslo, Norway
- 5 Department of Genitourinary Medicine, Guy's and St Thomas' NHS Foundation Trust, London, UK

## Address for Correspondence

Prof HJC de Vries, STI outpatient clinic, Cluster Infectious Diseases, Health Service Amsterdam, Amsterdam, The Netherlands. e-mail: <a href="mailto:h.j.devries@amc.uva.nl">h.j.devries@amc.uva.nl</a>

# WHAT IS NEW IN THIS UPDATED GUIDELINE?

# **Epidemiology**:

Based on clonal relatedness of prevalent LGV strains there is evidence that the LGV
epidemic among MSM in the Western world prevailed already in the United States in the
1980s and was introduced into Europe by the end of the last century.

# Aetiology and transmission

- A new LGV variant causing severe proctitis was unveiled and designated L2c.
- The L2b LGV variant causing the vast majority of infections among MSM is now also found among a few heterosexual women.

# Management:

- Apart from HIV and STI screening, HCV testing should be offered to all LGV patients.
- To exclude reinfections, STI screening during a follow up visit 3 months after an LGV diagnosis should be offered.

# LYMPHOGRANULOMA VENEREUM (LGV)

# **Epidemiology**

LGV has re-emerged among European men who have sex with men (MSM) in the past decade [1] and is probably endemic in this population where it is a relatively common cause of proctitis and occasional genital ulcer-adenopathy disease. LGV among MSM in Europe is caused in the majority of cases by the *C. trachomatis* biovar L2b which shows a high degree of clonal relatedness as found by Multi Locus Sequence Typing (MLST) in a molecular epidemiological study. [2] This is in contrast to the strains circulating among MSM in the United States, which show more molecular diversity. Based on these findings it is now speculated that the LGV epidemic among MSM in Europe caused by the L2b variant may have been imported to Europe from the United States by the end of the previous century via the highly internationalized network of sexual contacts among MSM. Whilst occasional cases of heterosexual LGV are seen in Europe these have usually been imported from endemic countries.

# **Aetiology and transmission**

- Causative pathogen: Chlamydia trachomatis types L1, L2, and L3. Additional variants have been described such as L2b, the strain currently found in MSM. A new LGV variant originating from a recombination of L2 and D C. trachomatis lineages has been reported but the extent of transmission of this hybrid organism within the community needs further investigation. [3] In contrast to serovars A-K which remain confined to the mucosa, serovar L strains are invasive organisms that disseminate via underlying connective tissue and spread to regional lymph nodes.
- Worldwide, LGV is thought to account for 2 to 10% of GUD in areas such as India and Africa. In Western regions LGV is endemic among MSM, mainly those co-infected with HIV.[ Heterosexual transmission of the LGV associated L2b strain has been described. [5, 6]
- Neither the degree of infectiousness nor the reservoir of disease has been accurately
  defined, but heterosexual transmission has been attributed largely to asymptomatic female
  carriers and in the MSM population, asymptomatic rectal infection and/or penile infection is
  the likely source of onward transmission.[7]

# **Clinical features**

Depending on the site of inoculation LGV can cause inguinal disease (usually after inoculation of the genitalia), or the anorectal syndrome (usually after inoculation via the rectum). The disease course usually follows three separate stages.

- Incubation period one to four weeks
- In the current LGV epidemic among MSM, proctitis is the primary manifestation of infection, usually presenting within a few weeks of sexual contact. It is characterised by severe symptoms of anorectal pain, haemopurulent discharge and bleeding per rectum; tenesmus and constipation are also seen due to the mucosal and perirectal oedema. Proctoscopic examination may reveal a distal granular or haemorrhagic proctitis with purulent exudate,

- mucosal ulceration and tumorous masses. LGV proctitis is notusually accompanied by inguinofemoral lymphadenopathy; however radiological imaging may demonstrate pelvic node involvement.[8]
- In the present LGV epidemic among MSM, symptomatology has varied in different populations, with UK cohorts showing almost all LGV to be symptomatic[9], in contrast to Dutch studies where a significant proportion of asymptomatic infection has been detected.[10, 11]
- LGV proctitis mimics chronic inflammatory bowel diseases like Crohn's disease, both clinically and in the pathological substrate.
- Primary Lesion small painless papule or pustule; may erode to form a small herpetiform ulcer. Usually heals within one week and often remains unnoticed. Mucopurulent discharge may be present, affecting the urethra, the cervix or the rectum depending on the inoculation site.
- Second stage: "inguinal stage" begins 2 to 6 weeks after onset of primary lesion. Causes painful inflammation of the inguinal and/or femoral lymph nodes. Typically this produces unilateral enlargement, inflammation, suppuration and abscesses. These "buboes" may become fluctuant and rupture in one third of patients. Some patients develop the "groove sign", which results from enlargement of the inguinal nodes above and the femoral nodes below Poupart's ligament.
- Inguino-femoral lymphadenopathy is mainly seen when the inoculation site is located on the
  external genitalia, which is the case in many male patients. In contrast, women more often
  have primary involvement of the rectum, upper vagina, cervix, or posterior urethra; as these
  regions drain to the deep iliac or perirectal nodes, inguinofemoral lymphadenopathy is not
  seen. The resultant intra-abdominal or retroperitoneal lymphadenopathy may lead to
  symptoms of lower abdominal pain or low back pain.
- Constitutional symptoms, such as low-grade fever, chills, malaise, myalgias and arthralgias
  may present during the second stage of disease. In addition, systemic spread of C.
  trachomatis occasionally results in arthritis, pneumonitis or (peri) hepatitis. Rare systemic
  complications include cardiac involvement, aseptic meningitis and ocular inflammatory
  disease.
- A rare presentation is the pharyngeal syndrome affecting the mouth and throat. Cervical lymphadenopathy and buboes can occur.[14]
- The third stage of disease in LGV is often called the "anogenitorectal syndrome" and is more often present in women. Patients initially develop proctocolitis followed by peri-rectal abscess, fistulas, strictures and stenosis of the rectum, possibly leading to "lymphorrhoids" (haemorrhoid-like swellings of obstructed rectal lymphatic tissue). Without treatment, chronic progressive lymphangitis leads to chronic oedema and sclerosing fibrosis, resulting in strictures and fistulas of the involved region, which can ultimately lead to elephantiasis, esthiomene (the chronic ulcerative disease of the external female genitalia) and the frozen

pelvis syndrome. If left untreated, LGV proctitis can lead to rectal stricture, with subsequent sequelae of soiling, pain, constipation and the possible development of mega colon.[15]

# **Diagnosis**

- The diagnosis of LGV is confirmed by the detection of biovar-specific *C. trachomatis* DNA in 1) ulcer material from primary anogenital lesions, 2) rectal specimens (in suspected cases of anorectal LGV); anorectal swabs are preferably collected from the mucosal lining under proctoscopic vision, alternatively a blind anorectal swab can suffice, or 3) bubo aspirates (in suspected cases inguinal LGV); historically LGV has been difficult to isolate from bubo aspirates in culture studies. (IV, C). Modern diagnostic approaches have been reviewed recently [11].
- Most modern laboratories follow a 2-step procedure:
  - First, a commercially available *C. trachomatis* NAAT test can be used to screen suspected samples. Although commercially available tests are not approved for extragenital sites, a large body of literature supports the use of these tests for the detection of rectal chlamydial infections.[11, 16-18] (III, B)
  - o If *C. trachomatis* is detected, LGV biovar-specific DNA then needs to be detected from the same specimen. For this purpose two "in house" NAAT tests have been reported; firstly, a real-time PCR-based test that specifically detects all *C. trachomatis* LGV biovar strains[19] and more recently, a real-time quadriplex PCR-based assay which incorporates both LGV-specific and non-LGV-specific target sequences, a *C. trachomatis* plasmid target, and the human RNase P gene as an internal control[20]. (III, B)
- If molecular diagnostic test facilities are not available, then a presumptive LGV diagnosis can be made using *Chlamydia* genus-specific serological assays. A high antibody titre (esp. IgA anti-MOMP antibodies) in a patient with a clinical syndrome suggestive of LGV supports the diagnosis.[21, 22] Nonetheless, a low titre does not exclude LGV, nor does a high titre in a patient without LGV symptomatology confirm LGV infection.[10,11] (III, B)The identification of rectal polymorphonuclear leucocytes (PMNLs) from rectal swabs is predictive of LGV proctitis, especially in HIV-positive MSM, with levels of >10 [11].

## Management

- It is recommended to screen all MSM who report receptive anal sexual practices in the previous 6 months for anorectal *C. trachomatis* infection. Subsequently, MSM who are anorectal *C. trachomatis* positive are then screened for LGV proctitis according to local guidelines.[11] (IIa)
- The prevalence of HIV among LGV cases ranges from 67% to 100% in 13 descriptive studies. There is a significant association between HIV and LGV (odds ratio 8.19, 95% CI 4.68-14.33). (4). (I, A)

 Tests for STI, including HIV (if not already known HIV-pos), hepatitis B and hepatitis C should be offered before starting therapy. (III, C) [23]

# Information, explanation and advice for the patient

- Patients should be informed that LGV is an invasive bacterial infection that is sexually transmitted but curable with antibiotics. Left untreated it can have serious and permanent adverse sequelae. Most of these complications are preventable if treatment is initiated in the early stages. (IV, C) Patient information leaflets focussed on the LGV epidemic among MSM are available from IUSTI (<a href="http://www.iusti.org/regions/europe/pdf/2010/PIL\_LGV.pdf">http://www.iusti.org/regions/europe/pdf/2010/PIL\_LGV.pdf</a>) and provided by several national organisations like the Terrence Higgins Fund (UK), and Schorer stichting (NL).
- Symptoms should resolve within 1-2 weeks of commencing antibiotic therapy. (III, B)
- Patients should abstain from any sexual contact until they have completed therapy. (IV, C)
- Screening for syphilis, C. trachomatis, N. gonorrhoeae, HIV, hepatitis B, and hepatitis C should be offered during a follow-up visit 3 months after an LGV diagnosis to exclude reinfections (IV, C) [23]

# **Therapy**

Despite a paucity of robust evidence[24] regarding the efficacy of therapy for any rectal chlamydial infections (LGV or non-LGV), three weeks of oral doxycycline 100mg twice daily to treat LGV is recommended [25-27]. The vast majority of recent MSM case reports have observed complete responses to this therapy; shorter courses may not eradicate the organism[28].

- First line doxycycline 100mg twice a day orally for 21 days. (IIb, B)
- Second line erythromycin 500mg four times a day orally for 21 days. (III, B)

\*Azithromycin in single- or multiple-dose regimens has also been proposed[23,29,30] but evidence is lacking to recommend this drug currently. (IV, C) Doxycycline is contraindicated in pregnancy and breastfeeding.

## Adjunctive therapy

- If fluctuant buboes appear they should be aspirated promptly through healthy adjacent skin. (IV, C)
- Surgical incision of buboes is not usually recommended due to potential complications such as chronic sinus formation. (IV, C)

 Patients with residual fibrotic lesions or fistulae do not benefit from further courses of antibiotics so surgical repair, including reconstructive genital surgery, should be considered.
 (IV, C)

# **Partner notification**

As LGV is sexually transmitted it is essential that partner notification is initiated when the diagnosis is made. Sexual contacts within the last 3 months should be offered testing for Chlamydia/LGV and empiric treatment with antibiotic therapy commenced until Chlamydia/LGV has been excluded in the partner. (IV, C)

#### Follow-up

All patients diagnosed with LGV should be followed up at the end of treatment:

- to ensure resolution of symptoms and signs of infection (IV, C)
- to check that adequate partner notification has been completed (IV, C)
- to address any patient concerns (IV, C)
- to arrange suitable follow-up testing for syphilis and blood-borne viruses including hepatitis
   B, C and HIV (IV, C)\*
- Although one case of doxycycline failure in LGV has been reported [31], a test of cure for LGV is not considered necessary if the recommended 21 day course of doxycycline is completed.[25] (III, B)

\*In the recent MSM LGV epidemic incident cases of both HIV and hepatitis C [28] have been observed and serological testing should be offered for both infections after appropriate window periods have elapsed according to relevant local guidelines.

# Prevention/health promotion

Patients diagnosed with LGV should be counselled regarding prevention of other STIs including HIV and hepatitis C:

- Offer regular sexual health screening including HIV testing
- Condom use should be demonstrated and promoted
- Offer Hepatitis A and B vaccination for MSM
- Patients at risk of HIV infection should be advised of the availability of post exposure prophylaxis for HIV
- In particular, HIV-positive MSM should be made aware of recent trends in hepatitis C epidemiology and warned of the risks of unprotected anal sex, serosorting, recreational drug use and mucosally-traumatic sexual practices such as fisting. Enema use may be implicated in LGV transmission [32]. Although sharing of equipment was rare, it is prudent to advise against sharing any such equipment and to wash equipment thoroughly after use.

## Auditable Outcome Measures (target 95% for all)

- All cases of suspected LGV should be subjected to laboratory investigations.
- All patients should be interviewed for the purpose of partner notification and this should be documented in the case notes. Sexual contacts within at least 3 months should be traced, tested and treated.
- HIV, syphilis, hepatitis B and hepatitis C serological testing as well as screening for concomitant other STIs, should be offered before starting therapy and after three months.
- Suspected or confirmed cases of LGV should be reported and relevant surveillance data collected according to local and national guidelines.

# Proposed date of revision May 2015

## **European STI Guidelines Editorial Board**

Dr Keith Radcliffe, UK - Editor-in-Chief

Dr Karen Babayan, Armenia (appointed 2009)

Dr Marco Cusini, Italy (app. 2010)

Dr Gilbert Donders, Belgium (app. 2012)

Prof Mikhail Gomberg, Russia (app. 2010)

Dr Michel Janier, France (app. 2006)

Dr Jorgen Skov Jensen, Denmark (app. 2006)

Prof. Harald Moi, Norway (app. 2007)

Dr Raj Patel, UK (app. 2006)

Prof Jonathan Ross, UK (app. 2006)

Dr Jackie Sherrard, UK (app. 2009)

Dr Magnus Unemo, Sweden (app. 2009)

Dr Willem van der Meijden, Netherlands (app. 2006)

Dr Simon Barton (UK) - UEMS representative, UK (app. 2010)

Dr Lali Khotenashvili – WHO European Office representative, Georgia (app. 2007)

Dr Marita van de Laar – ECDC representative, Netherlands (app. 2007)

Prof Mario Poljak - ESCMID representative, Slovenia (app. 2013)

Prof George-Sorin Tiplica, - EADV representative, Romania (app. 2012)

# List of contributing organisations

This guideline has been produced on behalf of the following organisations: the European Branch of the International Union against Sexually Transmitted Infections (IUSTI Europe); the European Academy of Dermatology and Venereology (EADV); the European Dermatology Forum (EDF); the European Society of Clinical Microbiology and Infectious Diseases (ESCMID); the Union of

European Medical Specialists (UEMS). The European Centre for Disease Prevention and Control (ECDC) and the European Office of the World Health Organisation (WHO-Europe) also contributed to its development.

- (1) White JA. Manifestations and management of lymphogranuloma venereum. Curr Opin Infect Dis 2009;22:57-66.
- (2) Christerson L, de Vries HJ, de Barbeyrac B, Gaydos CA, Henrich B, Hoffmann S, Schachter J, Thorvaldsen J, Vall-Mayans M, Klint M, Herrmann B, Morre SA. Typing of Lymphogranuloma Venereum Chlamydia trachomatis Strains. Emerg Infect Dis. 2010;**16**(11):1777-1779.
- (3) Somboonna N, Wan R, Ojcius DM, Pettengill MA, Joseph SJ, Chang A, Hsu R, Read TD, Dean D. Hypervirulent Chlamydia trachomatis clinical strain is a recombinant between lymphogranuloma venereum (L(2)) and D lineages. MBio. 2011;**2**(3):e00045-11.
- (4) Rönn MM, Ward H. The association between lymphogranuloma venereum and HIV among men who have sex with men: systematic review and meta-analysis. BMC Infect Dis. 2011;**11**:70.
- (5) Peuchant O, Baldit C, Le Roy C, Trombert-Paolantoni S, Clerc M, Bébéar C, de Barbeyrac B. First case of Chlamydia trachomatis L2b proctitis in a woman. Clin Microbiol Infect. 2011;**17**(12):E21-23.
- (6) Verweij SP, Ouburg S, de Vries H, Morré SA, van Ginkel CJ, Bos H, Sebens FW. The first case record of a female patient with bubonic lymphogranuloma venereum (LGV), serovariant L2b. Sex Transm Infect. 2012;**88**(5):346-347.
- (7) Vall-Mayans M, Caballero E, Sanz B. The emergence of lymphogranuloma venereum in Europe. Lancet 2009;**374**(9686):356.
- (8) van der Ham R, de Vries HJ. Lymphogranuloma venereum, where do we stand? Clinical recommendations. Drugs Today (Barc ) 2009;**45 Suppl B**:39-43.
- (9) Ward H, Martin I, Macdonald N, Alexander S, Simms I, Fenton K, French P, Dean G, Ison C. Lymphogranuloma venereum in the United kingdom. Clin Infect Dis 2007;**44**:26-32.
- (10) Spaargaren J, Fennema HS, Morré SA, de Vries HJ, Coutinho RA. New lymphogranuloma venereum Chlamydia trachomatis variant, Amsterdam. Emerg Infect Dis 2005;**11**(7):1090-1092.
- (11) van der Bij AK, Spaargaren J, Morre SA, Fennema HS, Mindel A, Coutinho RA, de Vries HJ. Diagnostic and clinical implications of anorectal lymphogranuloma venereum in men who have sex with men: a retrospective case-control study. Clin Infect Dis 2006;**42**:186-194.
- (12) Martin IM, Alexander SA, Ison CA, Macdonald N, McCarthy K, Ward H. Diagnosis of lymphogranuloma venereum from biopsy samples. Gut 2006;**55**:1522-1523.
- (13) Soni S, Srirajaskanthan R, Lucas SB, Alexander S, Wong T, White JA. Lymphogranuloma venereum proctitis masquerading as inflammatory bowel disease in 12 homosexual men. Aliment Pharmacol Ther. 2010;**32**(1):59-65.
- (14) Korhonen S, Hiltunen-Back E, Puolakkainen M. Genotyping of Chlamydia trachomatis in rectal and pharyngeal specimens: identification of LGV genotypes in Finland. Sex Transm Infect. 2012;88(6):465-469.
- (15) Pinsk I, Saloojee N, Friedlich M. Lymphogranuloma venereum as a cause of rectal stricture. Can J Surg 2007;**50**(6):E31-E32.
- (16) Halse TA, Musser KA, Limberger RJ. A multiplexed real-time PCR assay for rapid detection of Chlamydia trachomatis and identification of serovar L-2, the major cause of Lymphogranuloma venereum in New York. Mol Cell Probes 2006;**20**:290-297.
- (17) Hamill M, Benn P, Carder C, Copas A, Ward H, Ison C, French P. The clinical manifestations of anorectal infection with lymphogranuloma venereum (LGV) versus non-LGV strains of Chlamydia trachomatis: a case-control study in homosexual men. Int J STD AIDS 2007;18(7):472-475.
- (18) Jalal H, Stephen H, Alexander S, Carne C, Sonnex C. Development of real-time PCR assays for genotyping of Chlamydia trachomatis. J Clin Microbiol 2007;**45**:2649-2653.
- (19) Morré SA, Spaargaren J, Fennema JS, de Vries HJ, Coutinho RA, Peña AS. Real-time polymerase chain reaction to diagnose lymphogranuloma venereum. Emerg Infect Dis 2005; **11**(8):1311-1312.
- (20) Chen CY, Chi KH, Alexander S, Ison C, Ballard RC. A real-time quadriplex PCR assay for the diagnosis of rectal lymphogranuloma venereum (LGV) and non-LGV Chlamydia trachomatis infections. Sex Transm Infect 2008; 84(4):273-276.
- (21) de Vries HJ, Smelov V, Ouburg S, Pleijster J, Geskus RB, Speksnijder AG, Fennema JS, Morré SA. Anal lymphogranuloma venereum infection screening with IgA anti-Chlamydia trachomatis-specific major outer membrane protein serology. Sex Transm Dis. 2010;**37**(12):789-795.
- (22) van der Snoek EM, Ossewaarde JM, van der Meijden WI, Mulder PG, Thio HB. The use of serological titres of IgA and IgG in (early) discrimination between rectal infection with non-lymphogranuloma venereum and lymphogranuloma venereum serovars of Chlamydia trachomatis. Sex Transm Infect 2007;83(4):330-334.

- (23) Hill SC, Hodson L, Smith A. An audit on the management of lymphogranuloma venereum in a sexual health clinic in London, UK. Int J STD AIDS. 2010;**21**(11):772-776.
- (24) McLean CA, Stoner BP, Workowski KA. Treatment of lymphogranuloma venereum. Clin Infect Dis 2007;**44 Suppl 3**:S147-S152.
- (25) Centers for Disease Control and Prevention (CDC). Lymphogranuloma venereum among men who have sex with men--Netherlands, 2003-2004. MMWR Morb Mortal Wkly Rep 2004;**53**(42):985-988.
- (26) Clinical Effectiveness Group of the British Association for Sexual Health and HIV (CEG/BASHH). 2006 National Guideline for the Management of Lymphogranuloma Venereum (LGV) of the British Association of Sexual Health and HIV (BASHH). 2006.
- (27) Centers for Disease Control and Prevention (CDC), Workowski KA, Berman SM. Sexually transmitted diseases treatment guidelines, 2006. MMWR Recomm Rep 2006;55(RR-11):1-94.
- (28) de Vries HJ, Smelov V, Middelburg JG, Pleijster J, Speksnijder AG, Morré SA. Delayed microbial cure of lymphogranuloma venereum proctitis with doxycycline treatment. Clin Infect Dis 2009;48(5):e53-e56.
- (29) Nieuwenhuis RF, Ossewaarde JM, van der Meijden WI, Neumann HA. Unusual presentation of early lymphogranuloma venereum in an HIV-1 infected patient: effective treatment with 1 g azithromycin. Sex Transm Infect 2003;**79**(6):453-455.
- (30) van Nieuwkoop C, Gooskens J, Smit VT, Claas EC, van Hogezand RA, Kroes AC, Kroon FP. Lymphogranuloma venereum proctocolitis: mucosal T cell immunity of the rectum associated with chlamydial clearance and clinical recovery. Gut 2007; **56**(10):1476-1477.
- (31) Méchaï F, de Barbeyrac B, Aoun O, Mérens A, Imbert P, Rapp C. Doxycycline failure in lymphogranuloma venereum. Sex Transm Infect. 2010 Aug;86(4):278-279.
- (32) de Vries HJ, van der Bij AK, Fennema JS, Smit C, de Wolf F, Prins M, Coutinho RA, Morré SA. Lymphogranuloma venereum proctitis in men who have sex with men is associated with anal enema use and high-risk behavior. Sex Transm Dis 2008;**35**(2):203-208.

#### **APPENDICES**

# Search strategy:

MEDLINE and PubMed searches were performed from 2000 to December 2012 using Domain: HIV, MSM, homosexuals, men who have sex with men, Determinants: CT serovar L1-L3, CT infection\* L1-L3, Chlamydia L1-L3, Chlamydiasis L1-L3, LGV, lymphogranuloma venereum, Chlamydia trachomatis genovar L1-L3, Chlamydia trachomatis biovar L1-L3, Chlamydia trachomatis serovar L1-L3 Outcome: Anal infection, LGV, lymphogranuloma venereum

#### Levels of Evidence

la Evidence obtained from meta-analysis of randomised controlled trials.

**Ib** Evidence obtained from at least one randomised controlled trial.

**Ila** Evidence obtained from at least one well designed study without randomisation.

**IIb** Evidence obtained from at least one other type of well designed quasi-experimental study.

**III** Evidence obtained from well designed non-experimental descriptive studies such as comparative studies, correlation studies, and case control studies.

**IV** Evidence obtained from expert committee reports or opinions and/or clinical experience of respected authorities.

# **Grading of Recommendations**

A (Evidence levels Ia, Ib)

Requires at least one randomised control trial as part of the body of literature of overall good quality and consistency addressing the specific recommendation.

B (Evidence levels IIa, IIb, III)

Requires availability of well conducted clinical studies but no randomised clinical trials on the topic of recommendation.

C (Evidence IV)

Requires evidence from expert committee reports or opinions and/or clinical experience of respected authorities. Indicates absence of directly applicable studies of good quality.

## **Declarations of Interest**

Henry JC de Vries – none to declare Alexander Kreuter – none to declare Adele Zingoni - none to declare John A White – none to declare Harald Moi – none to declare