

2017 European guidelines for the management of genital herpes

Authors:

Rajul Patel¹, Oliver John Kennedy², Emily Clarke¹, Anna Geretti³, Arvid Nilsen⁴, Elizabeth Foley¹, Stephan Lautenschlager⁵, John Green⁶, Gilbert Donders⁷, Willem van der Meijden⁸, Mikhail Gomberg⁹, Harald Moi^{10,11} and Elizabeth Foley¹

1. Department of Genitourinary Medicine, Southampton, UK
2. Southampton Medical School, Southampton, UK
3. Department of Virology, University College London, London, UK
4. Department of Dermatovenerology, University of Bergen, Bergen, Norway
5. Dermatologisches Ambulatorium Triemli, Zurich, Switzerland
6. Central and North West London NHS Trust, London, UK
7. Department of Obstetrics and Gynecology, University Hospital Antwerp
8. Department of Dermatology, Erasmus Medical Centre, Rotterdam, The Netherlands
9. Department of Dermatology and Venereology, Moscow State University for Medicine and Dentistry, Moscow, Russia
10. Department of Venereology, the Olafia Clinic, Oslo University Hospital
11. Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Abstract:

Genital herpes is one of the commonest sexually transmitted infections worldwide. Using the best available evidence, this guideline recommends strategies for diagnosis, management and follow-up of the condition as well as for minimising transmission. Early recognition and initiation of therapy is key and may reduce the duration of illness or avoid hospitalisation with complications, including urinary retention, meningism or severe systemic illness. The guideline covers a range of common clinical scenarios, such as recurrent genital herpes, infection during pregnancy and coinfection with human immunodeficiency virus.

Declarations of interest:

None of the authors or editors had any conflicts of interest for the content of this guideline.

Keywords:

Herpes simplex, HSV, genital herpes, Europe, treatment, antibiotic

Revision date: 05/12/2016

INTRODUCTION

First infection with either herpes simplex virus type 1 (HSV-1) or type 2 (HSV-2) is termed primary infection and results in either symptomatic disease at the site of viral entry (i.e. on the face or genital area) or asymptomatic, and thus unrecognised, infection. In addition there may be systemic symptoms, as with other acute viral illnesses. Only a third of patients who acquire genital herpes have any recognised clinical features at the time of acquisition. Following infection, the virus becomes latent in

the local sensory ganglion, periodically reactivating to cause symptomatic lesions, or undergo asymptomatic, but nonetheless infectious, viral shedding. Genital herpes can be caused by either HSV-1 – the usual cause of oro-labial herpes – or by HSV-2. Infection with either virus can cause an identical initial illness; however, the actual clinical presentation may depend upon previous HSV-1 or HSV-2 infection, and previous sites of infection. Subsequent recurrence frequency is greater for HSV-2 than HSV-1 disease when infection involves the genital area.

Transmission risk

Risk of transmission appears to be greatest during lesional recurrences or prodrome, and patients should be advised to abstain from sexual contact during this time. Transmission can occur in the absence of lesional recurrence as a result of sub-clinical viral shedding. Efficacy of condoms to prevent sexual transmission has not been formally assessed; however, indirect evidence from failed vaccine trials provides strong support for their consistent usage (IIb, B).^{1,2}

DIAGNOSIS

Clinical diagnosis

Although classical genital herpes can be recognised by the presence of typical papular lesions progressing to vesicle and ulcer formation, associated with local adenitis and in recurrent cases preceded by prodromal symptoms, the features in many patients can be highly variable. The majority of patients will suffer from atypical lesions where signs may be easily confused with other genital dermatoses. In atypical cases, depending upon clinical diagnosis alone should be avoided, whenever possible.

Laboratory diagnosis

Virus detection

- Laboratory confirmation is recommended in all patients with suspected genital herpes, using methods that directly demonstrate the virus in genital specimens, typically swabs taken from the base of the lesion (vesicles should be unroofed with a needle or scalpel blade) and placed in viral transport medium (Ib, A).³
- HSV typing into HSV-1 and HSV-2 is recommended in all patients with first-episode genital herpes to guide counselling and management (III, B).¹
- As HSV shedding is intermittent, testing swabs from asymptomatic patients is not recommended for routine diagnosis since it is unlikely to yield confirmation of carrier status (Ib, A).¹
- HSV DNA detection is now considered the gold standard for diagnosis. It is both more sensitive and specific than HSV cell culture (Ib, A).³⁻⁸ Cell culture may occasionally be required to determine anti-viral sensitivity and to generate enough virus for transmission studies.
- HSV DNA detection by real-time PCR increases HSV detection rates in muco-cutaneous swabs by 11-71% compared with virus culture and is recommended as the preferred diagnostic method (Ib, A).^{3,7,8} Real-time PCR can tolerate less stringent conditions for sample storage and transport than virus culture, and allows the rapid detection and typing of HSV with a lower risk of contamination than traditional PCR assays.
- Viral antigen detection methods such as direct immunofluorescence assay (IFA), enzyme immunoassay (EIA) and Tzanck and Papanicolaou staining are no longer recommended except in extremely limited resource settings (Ib, A).⁹⁻¹¹

HSV type-specific serology

Serological testing is not routinely recommended in asymptomatic patients (IV, C), but may be useful in the following groups:^{1, 12-17}

- History of recurrent or atypical genital disease when direct virus detection methods have been negative (III, B). HSV-2 antibodies are supportive of a diagnosis of genital herpes; HSV-1 antibodies do not differentiate between genital and oro-pharyngeal infection. Counselling of HSV-2 IgG negative, HSV-1 IgG positive patients should take into account that HSV-1 is an uncommon cause of recurrent genital disease.¹
- First-episode genital herpes, where differentiating between primary and established infection guides counselling and management (III, B). At the onset of symptoms, the absence of HSV IgG against the virus type detected in the genital lesion is consistent with a primary infection.¹ Seroconversion should be demonstrated at follow-up.
- Sexual partners of patients with genital herpes, where concerns are raised about transmission. Serodiscordant couples can be counselled about strategies to reduce the risk of infection and disease (Ib, A).

Care must be taken in interpreting negative results since antibodies to both HSV-1 and 2 can be lost with time.

Testing of asymptomatic pregnant women is not routinely recommended, but is indicated when there is a history of genital herpes in the partner (IIb, B).¹⁸⁻²⁰ HSV-1 and/or HSV-2 seronegative women should be counselled about strategies to prevent a new infection with either virus type during pregnancy.

Limited data suggests an increased risk of perinatal HIV transmission among HSV-2 seropositive HIV-infected women.^{21, 22} As evidence is not consistent, testing of HIV positive pregnant women is not routinely recommended (IV, C).²³

HSV serological assays should be used that detect antibodies against the antigenically unique glycoproteins gG1 and gG2.^{11, 24} Non type-specific HSV antibody assays are of no value in the management of genital herpes.

Western blot (WB) is the diagnostic gold-standard. It is >97% sensitive and >98% specific, but is labour-intensive and not commercially available.^{25, 26}

The sensitivities and specificities of commercially available seroassays can vary significantly across populations.²⁷⁻³⁶ The positive predictive value (PPV) in low prevalence settings can make results uninterpretable. False negative results are more likely to occur in early infection and can be resolved by repeat testing.

HSV seroprevalence rates, the presence of risk factors for genital herpes, and clinical history influence the positive predictive value (PPV) of HSV type-specific serology and should guide testing and result interpretation (III, B).^{13, 22-31} The specificities of ELISA tests can be improved by raising the index value for interpreting positivity and testing all intermediate specimens with an alternate confirmatory test (IIa, B).^{26, 27, 29-31}

IgM testing is not recommended in routine clinical practice. Studies suggest that 12-30% of patients lose their HSV type specific antibodies depending on their HSV types and the test used.³⁷

MANAGEMENT

First-episode genital herpes

Indications for therapy

First episodes of genital herpes are frequently associated with a prolonged disease course. Untreated, many patients suffer general and local complications. Therapy can be highly effective and should be instigated at the earliest opportunity and on clinical suspicion alone.

Antivirals

Patients presenting within 5 days of the start of the episode, or while new lesions are still forming, should be given oral antiviral drugs. Aciclovir, valaciclovir and famciclovir are all effective in reducing the severity and duration of episode (Ib, A).^{38, 39} No therapy alters the natural course of genital herpes infection.

Topical agents are not recommended as they are less effective than oral agents and easily generate resistance (IV, C).⁴⁰

The only indication for the use of intravenous therapy is when the patient is unable to swallow or tolerate oral medication because of vomiting. The recommended regimens – all for five to ten days – are:

- Aciclovir 400 mg three times a day, or
- Aciclovir 200 mg five times a day, or
- Famciclovir 250 mg three times a day, or
- Valaciclovir 500 mg two times a day

Choice should be made by individual clinicians, taking cost of therapy and likely compliance into account. A number of patients will have extended episodes beyond five days. If a decision to provide five days of therapy is made, the patient should have early review to ensure they have no further lesion development, systemic symptoms or disease complications – they all will require extended therapy.

Supportive measures

Saline bathing and the use of appropriate analgesia are recommended. Although the potential for sensitisation exists in the use of topical anaesthetic agents, lignocaine is a rare sensitiser and can be used safely in genital herpes in the form of gel or ointment.⁴¹ Benzocaine, however, is a potent sensitiser and should not be used (IV, C).

Counselling

It is important to be frank about transmission risks including subclinical shedding and the limited impact of condoms and antivirals. Advice on disclosure should be practical and tailored to the patient's personal situation. The low physical morbidity and high population prevalence should be stressed. Clear information about pregnancy is important both to men and women. High distress at diagnosis is common, often persists with recurrences and may be reduced by antivirals (Ib, A).⁴²⁻⁴⁴ Most patients require one or two sessions but adjustment is difficult to predict and careful follow up is important with more intensive input for those who do not adjust within 3-6 months.

Management of complications

Hospitalisation may be required for urinary retention, meningism, severe constitutional symptoms, or adverse social circumstances. If catheterisation is required, consideration should be given as to whether a suprapubic approach offers better symptom control to the individual patient. Super infection of lesions is rare, but may occur during the second week. This is characterised by the recrudescence of local symptoms. Candida is most often implicated and is easily diagnosed and treated.

Special situations – HIV-positive patients with first-episode genital herpes

There are no controlled trials on duration and dose of treatment. Some clinicians advocate a 10-day course of treatment at twice the standard dose of any of the usual agents (IV, C).

Information for patients

The following information should be discussed when counselling patients with first episode genital herpes:

- The course of infection, including subclinical shedding
- Treatment options
- The risk of transmission, and interventions that may limit or reduce the risk of transmission
- The risk of transmission to the infant at birth. The patient should be counselled to inform the obstetrician or midwife
- The possibility of partner notification and the possible source of infection
- That transmissions can occur from an asymptomatic partner some years into a monogamous relationship

Follow-up

Patients are followed-up until the episode has resolved and counselling is considered complete. Further follow-up may be required to exclude other causes of genital ulceration that may be co-existent. Patients should be invited to re-attend should recurrences be problematic.

Recurrent genital herpes

Indications for therapy

Genital herpes recurrences are self-limiting and generally cause minor symptoms. Decisions about how best to manage clinical recurrences should be made in partnership with the patient. Management strategies include supportive therapy only, episodic antiviral treatments, and suppressive antiviral therapy. The most appropriate strategy for managing an individual patient may vary over time according to recurrence frequency, symptom severity, and relationship status. For most patients, management will need to be supportive only, with simple local measures such as saline bathing or topical petroleum jelly being adequate.

Episodic antiviral treatment

Oral aciclovir, valaciclovir and famciclovir are effective at reducing the duration and severity of recurrent genital herpes. The reduction in duration is a median of 1-2 days (Ib, A).⁴⁵⁻⁴⁷ Head-to-head studies of their effects show no advantage of one therapy over another, or the advantage of extended 5-day treatment over ultra-short therapy. Prodrugs offer simplified twice-a-day dosing. Patient-initiated

treatment started within 24 hours is most likely to be effective. Aborted lesions have been documented in up to one third of patients with early treatment.⁴⁸ To ensure prompt treatment, patients should be advised to carry a small quantity of drugs at all times.

Short course therapies should be tried in the first instance:

- Aciclovir 800 mg three times daily for 2 days, or⁴⁹
- Famciclovir 1 gram twice daily for one day, or⁵⁰
- Valaciclovir 500 mg twice daily for 3 days (Ib, A).^{49, 51-54}

Alternative longer 5 day courses include :

- Aciclovir 400 mg three times daily for 3-5 days, or
- Aciclovir 200 mg five times daily, or
- Valaciclovir 500 mg twice daily or
- Famciclovir 125 mg twice daily.

Suppressive therapy

The majority of early trials of suppressive therapy were done in patients with a recurrence rate equivalent to ≥ 6 recurrences/annum. More recently, studies have been completed in patients with milder disease including those with only serological evidence of infection. These indicate that patients across all spectrums of disease will benefit from a reduced rate of recurrence with treatment. The frequency of recurrence at which it is worth starting suppressive therapy is a subjective issue and needs to balance the frequency of recurrence, the impact of disease on the individual and the need to manage transmission risk against the cost and inconvenience of treatment.

All patients are highly likely to experience a substantial reduction in recurrence frequency on suppressive antiviral therapy. However, the majority of patients on such a regimen will still experience an occasional symptomatic recurrence.

Experience with suppressive antiviral therapy is most extensive with aciclovir (Ib, A).⁵⁵ Safety and resistance data on patients on long-term therapy now extends to over 18 years of continuous surveillance. Although not essential, it may be prudent to regularly assess the need for continuing therapy, since patient circumstances may alter significantly. However, even after prolonged periods of suppression, many patients do not find a significant alteration in disease frequency or severity upon discontinuation and reassessment.

Recommended regimens

The optimal total daily dose of suppressive aciclovir therapy is 800 mg. The only published clinical dose-ranging study concluded that 200 mg 4 times a day was marginally superior to 400 mg twice daily ($p < 0.02$) (Iib, B).⁵⁶ However, ability to comply with a 4 times a day dosing regimen should determine prescribing decisions for individual patients. Twice-daily valaciclovir (250 mg twice daily) has been shown to be as effective as twice-daily aciclovir (400 mg twice daily). Once-daily aciclovir does not suppress genital herpes recurrences. There is some debate as to whether once daily therapy is as effective as twice daily therapy with valaciclovir. For those patients experiencing < 10 recurrences per annum, a dose of 500 mg daily valaciclovir will be adequate; for those patients experiencing > 10 recurrences per annum, 250 mg b.d. or 1 g once a day is required.⁵⁷

No major clinically significant differences between suppressive therapy with valaciclovir (500 mg daily) and famciclovir (250 mg twice daily) have been documented (IV, C).⁸ In patients with an insufficient clinical response the daily suppressive dose of valaciclovir or famciclovir may have to be doubled (IV, C). Routine blood monitoring of standard dose therapy is not required. Occasionally a mild headache or nausea may occur with valaciclovir.

The decision to continue suppressive therapy should be reviewed at least annually. Discontinuation of therapy at this time, if the patient is willing, will allow a reassessment of recurrence frequency. A small number of patients will experience a reduction in recurrence frequency compared with pre-suppression symptomatic levels. The minimum period of assessment should include two recurrences to allow a view to be taken both on the frequency and severity. It is safe and reasonable to restart treatment in patients who continue to have significant disease (IV, C).

Short courses of suppressive therapy to prevent clinical symptoms may be helpful for some patients (e.g. for holidays, exams, etc.). Clinicians need to note that full suppressive effect is usually only obtained five days into treatment.

Viral shedding and transmission on suppressive therapy

Subclinical shedding of infectious virus occurs in most individuals with genital HSV-1 and/or HSV-2. Viral shedding is more likely to occur in patients with genital HSV-2, in the first year after infection, or in individuals with frequent symptomatic recurrences. Aciclovir, valaciclovir and famciclovir all suppress symptomatic and asymptomatic viral shedding.

Even if it seems biologically plausible, partial suppression of viral shedding does not necessarily equate to reduced transmission. However, suppressive therapy with valaciclovir 500 mg a day (in those with 10 or fewer recurrent episodes per year) significantly reduced transmission – by nearly 50% - in serodiscordant couples (Ib, A).¹⁶ Twice daily aciclovir 400mg achieves similar levels of reduction in asymptomatic shedding to once daily VACV. Suppressive antiviral therapy may be considered in addition to the use of condoms and selective sexual abstinence when transmission concerns are present.

SPECIAL SITUATIONS

Management of HSV in the immunocompromised and HIV positive patient

There is epidemiological synergy between Herpes simplex virus (HSV) and HIV infections.^{58, 59} Herpes simplex infections activate HIV replication and may facilitate onward HIV transmission to sexual partners.⁶⁰⁻⁶⁷ Suppressive treatment of HSV-2 infection with valaciclovir has been shown to reduce genital HIV shedding in women (not on ARVs).⁶⁸ In addition, both prevalent and incident HSV-2 infections are associated with an increased risk of HIV acquisition.^{69, 70}

The natural history of genital herpes in untreated people with HIV (PWHIV) is significantly different from that in HIV-negative individuals. The most important risk factor for herpes reactivation is the degree of HIV associated immunosuppression.⁷¹⁻⁷³

Standard systemic antiviral drugs, as used to treat genital herpes in HIV-uninfected patients, have been shown to successfully treat genital herpes in PWHIV.⁷⁴⁻⁷⁹ Resistance to anti-herpes drugs is more common in those with HIV co-infection and is associated with treatment failure of genital herpes.⁸⁰ Suppressive antiviral therapy with currently available agents has been shown in multiple studies to have

no impact on HIV acquisition or transmission risk. HSV treatment used only to manage or reduce HIV transmission or acquisition risk cannot be recommended (1b, A).^{81, 82}

Much of the evidence on herpes management in PWHIV comes from studies performed before the era of combination antiretroviral therapy; prospective studies performed early in the epidemic showed that clinical lesions might be persistent and progressive in those with HIV. Genital herpes, including chronic erosive lesions may occur as a manifestation of the immune reconstitution inflammatory syndrome (IRIS) following combination antiretroviral therapy.⁸³⁻⁸⁷ HSV associated IRIS may be unresponsive to previously effective anti-herpes viral therapy in the absence of increased antiviral resistance. Management is difficult but topical cidofovir may be effective.⁸⁸

Management of initial episode HSV

There is no trial data for any antiviral in initial episode genital HSV in HIV infected patients. The vast majority of adults with HIV have serological evidence of established HSV-1 and -2 infections making acquisition trials extremely difficult to perform.

Case studies report that acquisition of genital HSV may be associated with a prolonged and uncertain clinical course. Systemic symptoms may predominate and chronic lesions may become established if immunological clearance of the skin does not occur. In the absence of HIV therapy, primary genital herpes may be severe and prolonged with risk of progressive, multifocal, and coalescing mucocutaneous anogenital lesions. Moreover, serious and potentially life-threatening systemic complications such as fulminant hepatitis, pneumonia, neurological disease and disseminated infection, have been reported.

In the absence of data most authorities advise that multiples of the standard levels of treatment for first episode HSV be used in the immunocompromised. However, for those with HIV these may not always be required particularly for those with normal CD4 counts.

In patients with advanced HIV double the standard dose of antiviral should be considered and if new lesions continue to form at day 3-5 a higher dose should be considered. Prompt initiation of therapy is recommended. If new lesions are still forming after 3-5 days, a repeat viral isolation should be attempted and susceptibility testing arranged if possible. The dose of HSV therapy should also be increased.

Recommended initial doses:⁸⁹

- Aciclovir 400mg five times daily, for 7-10 days (IV,C)
- Valaciclovir 500-1g twice daily, for 10 days (IV,C)
- Famciclovir 250-500mg three times daily, for 10 days (IV,C)

Treatment should be given for at least 10 days or until all lesions have re-epithelialized – this will often exceed the usual 10 day duration of treatment that is given to HIV negative patients.

If fulminant disease ensues, then intravenous aciclovir be substituted at 5-10mg/kg body weight every 8 hours, for 2-7 days or until clinical improvement, and followed by oral antiviral therapy to complete a minimum of 10 days total treatment (IV, C).

Management of recurrent disease

Both clinical and subclinical reactivations of genital herpes are more frequent in people with HIV and may lead to persistent and progressive anogenital mucocutaneous lesions, especially with CD4 cell counts <50 per mm^3 . Features can be atypical in nature and hypertrophic lesions can occur. Optimising the control of HIV replication with combination antiretroviral therapy is of fundamental importance for the management of recurrent genital herpes. HAART will reduce the frequency of clinical recurrences but has less effect upon asymptomatic viral shedding. Thereafter, specific antiviral drugs can be used for either episodic or suppressive treatment. A number of trials of antiviral therapy in the immunocompromised have been reported.

Episodic therapy

It is likely that 5 days of therapy will be adequate for most patients. It should be noted that with advanced HIV 13-17% of patients have been reported to have new lesions developing at the end of a seven day course of treatment. Shorter courses of therapy may be adequate in those with good CD4 counts (>500 cell/ mm^3) although only one trial with famciclovir has reported this effect (1b,B).⁵⁰

Standard doses of antivirals should suffice in those with no evidence of immune failure (1b,A). In those with advanced disease it may be necessary to double the standard dose and to continue therapy beyond 5 days (1b,B). Caution should be exercised in using ultrashort courses of episodic therapy since these have not been evaluated fully in the immunocompromised.

Suppressive therapy

Suppressive antiviral therapy for HSV appears to be less effective in people with HIV than in HIV negative people, but remains well tolerated. All three agents have been trialled. Standard suppressive doses of aciclovir are effective. Valaciclovir is more effective when given twice daily (500mg bid) compared to once daily dosing (1000mg). The valaciclovir 500mg once daily dose has not been evaluated in the HIV positive patient. Trial data for the efficacy of high dose famciclovir is only available over much shorter durations.

It is recommended that intermittent cessation of suppressive antiviral therapy for genital herpes should occur, especially in those in whom there is also adequate inhibition of HIV replication and rising CD4 cell counts. In some PWHIV with less frequent outbreaks of genital herpes, episodic treatment may be substituted. In others, where the pre-treatment pattern of recurrences resumes, suppressive treatment may need to restart (IV, C).

There is a considerable body of data on the safety of oral antivirals in the HIV positive immunocompromised host. Two studies in the pre-HAART era looked at high dose aciclovir (400mg qid) and more recently at standard dose regimens. For valaciclovir a number of studies looked at the value of valaciclovir for the suppression of recurrent genital herpes. High dose valaciclovir (2g qid) has been studied and reported in HIV positive people, and those immunosuppressed and recovering from bone marrow transplants. Most recently a large number of studies looking at the efficacy of aciclovir and valaciclovir suppression and its impact on HIV transmission from co-infected patients have been reported. These trials indicate that use of oral aciclovir at standard dose and valaciclovir at 1g od and 500mg bd is associated with little or no adverse effect or toxicity as compared to the non HIV positive. High dose valaciclovir (8g daily) has been associated with Microangiopathic Haemolytic Uraemic syndrome.

Recommended drug regimens for daily suppressive treatment:⁸⁹⁻⁹¹

- Aciclovir 400 mg orally twice to three times a day
- Valaciclovir 500 mg orally twice a day.

If these options do not adequately control disease then the first option should be to double the dose. If control is still not achieved then famciclovir 500 mg orally twice a day can be tried (IIa,B).

HSV suppression to limit HIV progression

Suppressive antiviral therapy with aciclovir or valaciclovir has been shown to decrease the levels of HIV viraemia in those patients with detectable HIV viral loads through a mechanism not yet fully elucidated.⁹² Such a strategy will impact on HIV progression, particularly for those individuals not on HAART. A large RCT in early HIV (those individuals not on HAART and with CD4 counts above 250 cells/mm³) has shown that standard doses of suppressive antiviral therapy (aciclovir 400mg bd) will sustain CD4 counts above 250 cell/mm³ and this effect reduced the need for HAART at 2 years by 16% in the treatment group. However, benefits of suppressive antivirals have not been demonstrated in the presence of effective HAART.^{81, 82}


Management of recalcitrant herpes in immunocompromised individuals

Although rare in immunocompetent individuals, clinically refractory lesions due to genital HSV are a major problem in patients with severe immunodeficiency, including late-stage HIV diseases and patients with immune reconstitution inflammatory syndrome (IRIS) following combination antiretroviral therapy. Algorithms for treatment in such situations are shown in Figure 1. Systemic therapy with either foscarnet or cidofovir is generally preferred to treat drug resistant herpes in those with HIV. There is evidence for alternating courses of treatment with aciclovir and cidofovir for subsequent recurrences as a strategy that may reduce the development of cidofovir-resistant strains. The efficacy, safety, and durability of the therapeutic response of these agents have yet to be determined in prospective controlled trials.

In prospective studies, aciclovir-resistant strains have been found in around 5-7% isolates from genital herpes lesions in HIV-infected persons.^{40, 93, 94} Aciclovir resistance is confirmed if isolates require aciclovir concentrations >1-3 mg/l for inhibition. Aciclovir resistance is most commonly related to a mutation in the gene encoding HSV thymidine kinase (TK), which is responsible for initial phosphorylation of aciclovir to its active form, resulting in TK that either has reduced affinity for aciclovir or is not synthesised. TK-deficient strains are of reduced pathogenicity in immunocompetent individuals but may cause serious local and systemic disease in severely immunocompromised individuals.^{95, 96} They appear less likely to be associated with the development of latency; hence, subsequent clinical reactivations of genital herpes are often caused by aciclovir sensitive isolates. Partially resistant strains may sometimes be successfully treated with high dose intravenous aciclovir and other nucleoside analogues but fully aciclovir-resistant strains are resistant to valaciclovir and ganciclovir, and the majority are resistant to famciclovir.⁹⁵⁻⁹⁷ TK-deficient strains are susceptible to foscarnet and cidofovir which do not depend upon TK but which inhibit viral DNA polymerase.

Antiviral susceptibility testing for HSV has limited availability and therefore the clinical response to antiviral therapy is often used to guide decisions, Advice from a clinical virologist about appropriate drug dosages and duration may be sought when clinical resistance is suspected.

Figure 1: Algorithm for the treatment of herpes in immunocompromised individuals.



*All HSV strains resistant to aciclovir are also resistant to valaciclovir

** These products will often have to be made up in house

Management of partners

There is no evidence on which to base recommendations for partner notification. On an individual basis, it may be appropriate to offer to see partners to help with the counselling process. Partner notification in relation to pregnancy is discussed below. It is worth considering the following points when counselling partners;

- The use of condoms is advisable especially when transmission concerns are present - even when the index case is on suppressive antiviral treatment.¹⁶
- Asymptomatic shedding plays a major role in the transmission of HSV infection and selective abstinence in itself is not an effective strategy for managing transmission risk.

- Partner notification is an effective way of detecting uninfected or asymptomatic individuals especially when combined with type-specific antibody testing.
- Up to 50% of asymptomatic HSV-2 seropositive women can be taught to recognise genital herpes recurrences after counselling.
- Virus transmission can be reduced either with suppressive antiviral treatment or by using condoms.

Management of pregnant women with first episode genital herpes

First and second trimester acquisition

Management of the woman should be in line with her clinical condition and will often involve the use of either oral or intravenous aciclovir in standard doses.

Providing that delivery does not ensue, the pregnancy should be managed expectantly and vaginal delivery anticipated (IV, C).

Daily suppressive aciclovir 400mg t.i.d. from 36 weeks gestation may prevent HSV lesions at term and hence the need for delivery by Caesarean section (Ib, B).⁹⁸⁻¹⁰³

Third trimester acquisition (IV, C)

Caesarean section should be considered for all women, particularly those developing symptoms within 6 weeks of delivery, as the risk of viral shedding in labour is very high (Ib, B).

Daily suppressive aciclovir 400mg t.i.d. to delivery should be considered.

If vaginal delivery is unavoidable, prolonged rupture of membranes and invasive procedures, including the use of scalp electrodes, should be avoided. Intrapartum IV aciclovir given to the mother and subsequently to the baby may be considered and the paediatrician should be informed.¹⁰⁴

Management of pregnant women with recurrent genital herpes (III, B)

Women with recurrent genital herpes should be informed that the risk of neonatal herpes is low.

Symptomatic recurrences of genital herpes during the third trimester will be brief; vaginal delivery is appropriate if no lesions are present at delivery.

For women with a history of recurrent genital herpes who would opt for a Caesarean section if they had HSV lesions at the onset of labour, daily suppressive aciclovir 400mg tid from 36 weeks gestation may prevent HSV lesions at term and hence the need for delivery by Caesarean section (Ia, A).¹⁰⁵

If there are no genital lesions at delivery, there is no indication for a Caesarean section to prevent neonatal herpes.

Sequential cultures or PCR during late gestation to predict viral shedding at term are not indicated.¹⁰⁶ The utility of taking cultures or PCR at delivery, in order to identify women who are asymptotically shedding HSV, is unproven.

Management of recurrent HSV in early pregnancy

Although the safety of aciclovir in early pregnancy is not fully established judicious use of this agent for suspected acquisition episodes is widely advocated. The same cannot be said for recurrent disease. Continuous or episodic therapy is not recommended in early pregnancy and should be avoided. Clinicians are on occasion obliged to use therapy for severe and complicated disease and a case-by-case assessment should be made. Newer antivirals should be avoided and the dose of aciclovir titrated down to the minimum effective level.

Management of HIV positive women with recurrent HSV infection (IV, C)

There is some evidence that HIV antibody positive women with genital HSV ulceration in pregnancy are more likely to transmit HIV infection independent of other factors.^{21, 23} However, this is not a consistent finding across all studies.²³

Women who are HIV antibody positive and have a history of genital herpes should be offered daily suppressive aciclovir 400mg t.i.d. from 32 weeks gestation to reduce the risk of transmission of HIV-1 infection especially in women where a vaginal delivery is planned. Starting therapy at an earlier gestation than usual should be considered in view of the increased possibility of preterm labour (IV, C). There is currently no evidence to recommend daily suppressive treatment for HIV-1 antibody positive women who are HSV 1 or 2 seropositive but have no history of genital herpes.¹⁰⁷

Management of women with genital lesions at onset of labour

Caesarean section may be considered for women with recurrent genital herpes lesions at the onset of labour but the risk of neonatal herpes following vaginal delivery is small and must be set against risks to the mother of Caesarean section. Evidence from the Netherlands shows that a conservative approach, allowing vaginal delivery in the presence of an anogenital lesion, has not been associated with a rise in numbers of neonatal HSV cases (III, B).¹⁰⁸ However, this approach can only be adopted if fully supported by obstetricians and neonatologists, and if consistent with local medico-legal advice.

Clinical diagnosis of genital herpes at the time of labour correlates relatively poorly with HSV detection from genital sites by either culture or PCR and fails to identify women with asymptomatic HSV shedding.

Note: None of the antiviral drugs is licensed for use in pregnancy but the use of aciclovir in pregnancy has not been associated with any consistent pregnancy or foetal/neonatal adverse effects other than transient neutropenia.^{108, 109} Safety data for aciclovir may be extrapolated to valaciclovir in late pregnancy, as it is the valine ester, but there is less experience with use of valaciclovir.¹¹⁰ Famciclovir should currently be avoided.

Prevention of acquisition of infection (IV, C)

Maternal risk of HSV acquisition in pregnancy varies geographically and local epidemiological surveillance should guide strategy for prevention. Any strategy for prevention of neonatal herpes needs to involve both parents.

All women should be asked at their first antenatal visit if they or their partner have had genital or oral herpes.

Female partners of men with genital herpes, but without a personal history of genital herpes should be advised about reducing their risk of acquiring herpes in pregnancy and of subsequent transmission to

their baby. Strategies include selective and complete abstinence (especially in the third trimester) and conscientious condom use.

Daily suppressive treatment has been shown to significantly reduce the risk of transmission of HSV to a seronegative partner, however the effectiveness of suppressive treatment of the male partner to reduce transmission to a pregnant woman has not been evaluated so can currently only be recommended with caveats.

Pregnant women should be advised of the risk of acquiring HSV-1 as a result of receptive orogenital contact especially in the last trimester of pregnancy.

Identifying susceptible women by means of type-specific antibody testing has not been shown to be cost effective and is not indicated in Europe.

All women, not just those with a history of genital herpes, should undergo careful vulval inspection at the onset of labour to look for clinical signs of herpes infection.

Mothers, staff and other relatives/friends with active oral HSV lesions or herpetic whitlow should be advised to avoid direct contact between lesions and the neonate.

Management of the neonate

Babies born to mothers with first-episode genital herpes at the onset of labour

The paediatrician should be informed.

HSV PCR of urine and stool, from the oropharynx, eyes and surface sites, should be taken to allow early identification of infected babies

The potential benefits and risks of starting intravenous aciclovir without waiting for the results of these cultures should be discussed

If aciclovir is not started immediately the neonate should be closely monitored for signs of lethargy, fever, poor feeding or lesions.

Babies born to mothers with recurrent genital herpes at the onset of labour

Although some clinicians feel that taking a set of specimens for viral culture collected after delivery may help with early identification of infection there is no evidence to support this practice. However health care workers and parents should be advised to consider HSV in the differential diagnosis if the baby shows any signs of infection or develops skin, eye or mucous membrane lesions, particularly in the first two weeks of life.

Proposed date for review: 2022

Acknowledgements:

Composition of editorial board:

Please refer to document at: http://www.iusti.org/regions/Europe/pdf/2014/Editorial_Board2014.pdf

List of contributing organisations:

Please refer to text at: <http://www.iusti.org/regions/Europe/euroguidelines.htm>

References

1. Casper C, Wald A. Condom use and the prevention of genital herpes acquisition. *Herpes*. 2002;9(1):10-4.
2. Wald A, Langenberg AG, Krantz E, Douglas JM, Jr., Handsfield HH, DiCarlo RP, et al. The relationship between condom use and herpes simplex virus acquisition. *Ann Intern Med*. 2005;143(10):707-13.
3. Gupta R, Warren T, Wald A. Genital herpes. *Lancet*. 2007;370(9605):2127-37.
4. Koutsky LA, Stevens CE, Holmes KK, Ashley RL, Kiviat NB, Critchlow CW, et al. Underdiagnosis of genital herpes by current clinical and viral-isolation procedures. *N Engl J Med*. 1992;326(23):1533-9.
5. Wald A, Huang ML, Carrell D, Selke S, Corey L. Polymerase chain reaction for detection of herpes simplex virus (HSV) DNA on mucosal surfaces: comparison with HSV isolation in cell culture. *J Infect Dis*. 2003;188(9):1345-51.
6. Ramaswamy M, McDonald C, Smith M, Thomas D, Maxwell S, Tenant-Flowers M, et al. Diagnosis of genital herpes by real time PCR in routine clinical practice. *Sex Transm Infect*. 2004;80(5):406-10.
7. van Doornum GJ, Guldemeester J, Osterhaus AD, Niesters HG. Diagnosing herpesvirus infections by real-time amplification and rapid culture. *J Clin Microbiol*. 2003;41(2):576-80.
8. Geretti AM. Genital herpes. *Sex Transm Infect*. 2006;82 Suppl 4:iv31-4.
9. Verano L, Michalski FJ. Herpes simplex virus antigen direct detection in standard virus transport medium by Du Pont HerpChek enzyme-linked immunosorbent assay. *J Clin Microbiol*. 1990;28(11):2555-8.
10. Slomka MJ, Emery L, Munday PE, Mouldsdale M, Brown DW. A comparison of PCR with virus isolation and direct antigen detection for diagnosis and typing of genital herpes. *J Med Virol*. 1998;55(2):177-83.
11. Cone RW, Swenson PD, Hobson AC, Remington M, Corey L. Herpes simplex virus detection from genital lesions: a comparative study using antigen detection (HerpChek) and culture. *J Clin Microbiol*. 1993;31(7):1774-6.
12. Munday PE, Vuddamalay J, Slomka MJ, Brown DW. Role of type specific herpes simplex virus serology in the diagnosis and management of genital herpes. *Sex Transm Infect*. 1998;74(3):175-8.
13. Ashley RL, Wald A. Genital herpes: review of the epidemic and potential use of type-specific serology. *Clin Microbiol Rev*. 1999;12(1):1-8.
14. Malkin JE. Herpes simplex virus: who should be tested? *Herpes*. 2002;9(2):31.
15. Copas AJ, Cowan FM, Cunningham AL, Mindel A. An evidence based approach to testing for antibody to herpes simplex virus type 2. *Sex Transm Infect*. 2002;78(6):430-4.
16. Corey L, Wald A, Patel R, Sacks SL, Tyring SK, Warren T, et al. Once-daily valacyclovir to reduce the risk of transmission of genital herpes. *N Engl J Med*. 2004;350(1):11-20.
17. Ramaswamy M, McDonald C, Sabin C, Tenant-Flowers M, Smith M, Geretti AM. The epidemiology of genital infection with herpes simplex virus types 1 and 2 in genitourinary medicine attendees in inner London. *Sex Transm Infect*. 2005;81(4):306-8.
18. Brown ZA, Selke S, Zeh J, Kopelman J, Maslow A, Ashley RL, et al. The acquisition of herpes simplex virus during pregnancy. *N Engl J Med*. 1997;337(8):509-15.
19. Rouse DJ, Stringer JS. An appraisal of screening for maternal type-specific herpes simplex virus antibodies to prevent neonatal herpes. *Am J Obstet Gynecol*. 2000;183(2):400-6.
20. Tita AT, Grobman WA, Rouse DJ. Antenatal herpes serologic screening: an appraisal of the evidence. *Obstet Gynecol*. 2006;108(5):1247-53.
21. Drake AL, John-Stewart GC, Wald A, Mbori-Ngacha DA, Bosire R, Wamalwa DC, et al. Herpes simplex virus type 2 and risk of intrapartum human immunodeficiency virus transmission. *Obstet Gynecol*. 2007;109(2 Pt 1):403-9.

22. Bollen LJ, Whitehead SJ, Mock PA, Leelawiwat W, Asavapiriyant S, Chalermchockchareonkit A, et al. Maternal herpes simplex virus type 2 coinfection increases the risk of perinatal HIV transmission: possibility to further decrease transmission? *AIDS*. 2008;22(10):1169-76.
23. Chen KT, Tuomala RE, Chu C, Huang ML, Watts DH, Zorrilla CD, et al. No association between antepartum serologic and genital tract evidence of herpes simplex virus-2 coinfection and perinatal HIV-1 transmission. *Am J Obstet Gynecol*. 2008;198(4):399 e1-5.
24. Ashley RL. Performance and use of HSV type-specific serology test kits. *Herpes*. 2002;9(2):38-45.
25. Ashley R, Benedetti J, Corey L. Humoral immune response to HSV-1 and HSV-2 viral proteins in patients with primary genital herpes. *J Med Virol*. 1985;17(2):153-66.
26. Ashley R. Type-specific antibodies to HSV-1 and-2: review of methodology. *Herpes*. 1998;5:33-8.
27. Smith JS, Bailey RC, Westreich DJ, Maclean I, Agot K, Ndinya-Achola JO, et al. Herpes simplex virus type 2 antibody detection performance in Kisumu, Kenya, using the Herpesselect ELISA, Kalon ELISA, Western blot and inhibition testing. *Sex Transm Infect*. 2009;85(2):92-6.
28. Gopal R, Gibbs T, Slomka MJ, Whitworth J, Carpenter LM, Vyse A, et al. A monoclonal blocking EIA for herpes simplex virus type 2 antibody: validation for seroepidemiological studies in Africa. *J Virol Methods*. 2000;87(1-2):71-80.
29. Morrow RA, Friedrich D, Krantz E. Performance of the focus and Kalon enzyme-linked immunosorbent assays for antibodies to herpes simplex virus type 2 glycoprotein G in culture-documented cases of genital herpes. *J Clin Microbiol*. 2003;41(11):5212-4.
30. van Dyck E, Buve A, Weiss HA, Glynn JR, Brown DW, De Deken B, et al. Performance of commercially available enzyme immunoassays for detection of antibodies against herpes simplex virus type 2 in African populations. *J Clin Microbiol*. 2004;42(7):2961-5.
31. Golden MR, Ashley-Morrow R, Swenson P, Hogrefe WR, Handsfield HH, Wald A. Herpes simplex virus type 2 (HSV-2) Western blot confirmatory testing among men testing positive for HSV-2 using the focus enzyme-linked immunosorbent assay in a sexually transmitted disease clinic. *Sex Transm Dis*. 2005;32(12):771-7.
32. Morrow RA, Friedrich D, Meier A, Corey L. Use of "biokit HSV-2 Rapid Assay" to improve the positive predictive value of Focus HerpeSelect HSV-2 ELISA. *BMC Infect Dis*. 2005;5:84.
33. Ashley Morrow R, Krantz E, Friedrich D, Wald A. Clinical correlates of index values in the focus HerpeSelect ELISA for antibodies to herpes simplex virus type 2 (HSV-2). *J Clin Virol*. 2006;36(2):141-5.
34. Nascimento MC, Ferreira S, Sabino E, Hamilton I, Parry J, Pannuti CS, et al. Performance of the HerpeSelect (Focus) and Kalon enzyme-linked immunosorbent assays for detection of antibodies against herpes simplex virus type 2 by use of monoclonal antibody-blocking enzyme immunoassay and clinicovirological reference standards in Brazil. *J Clin Microbiol*. 2007;45(7):2309-11.
35. LeGoff J, Mayaud P, Gresenguet G, Weiss HA, Nzambi K, Frost E, et al. Performance of HerpeSelect and Kalon assays in detection of antibodies to herpes simplex virus type 2. *J Clin Microbiol*. 2008;46(6):1914-8.
36. Gamiel JL, Tobian AA, Laeyendecker OB, Reynolds SJ, Morrow RA, Serwadda D, et al. Improved performance of enzyme-linked immunosorbent assays and the effect of human immunodeficiency virus coinfection on the serologic detection of herpes simplex virus type 2 in Rakai, Uganda. *Clin Vaccine Immunol*. 2008;15(5):888-90.
37. van Rooijen MS, Roest W, Hansen G, Kwa D, de Vries HJ. False-negative type-specific glycoprotein G antibody responses in STI clinic patients with recurrent HSV-1 or HSV-2 DNA positive genital herpes, The Netherlands. *Sex Transm Infect*. 2016;92(4):257-60.
38. Corey L, Benedetti J, Critchlow C, Mertz G, Douglas J, Fife K, et al. Treatment of primary first-episode genital herpes simplex virus infections with acyclovir: results of topical, intravenous and oral therapy. *J Antimicrob Chemother*. 1983;12 Suppl B:79-88.

39. Fife KH, Barbarash RA, Rudolph T, Degregorio B, Roth R. Valaciclovir versus acyclovir in the treatment of first-episode genital herpes infection. Results of an international, multicenter, double-blind, randomized clinical trial. The Valaciclovir International Herpes Simplex Virus Study Group. *Sex Transm Dis.* 1997;24(8):481-6.
40. Reyes M, Shaik NS, Graber JM, Nisenbaum R, Wetherall NT, Fukuda K, et al. Acyclovir-resistant genital herpes among persons attending sexually transmitted disease and human immunodeficiency virus clinics. *Arch Intern Med.* 2003;163(1):76-80.
41. Weightman W, Turner T. Allergic contact dermatitis from lignocaine: report of 29 cases and review of the literature. *Contact Dermatitis.* 1998;39(5):265-6.
42. Carney O, Ross E, Bunker C, Ikkos G, Mindel A. A prospective study of the psychological impact on patients with a first episode of genital herpes. *Genitourin Med.* 1994;70(1):40-5.
43. Patel R, Tyring S, Strand A, Price MJ, Grant DM. Impact of suppressive antiviral therapy on the health related quality of life of patients with recurrent genital herpes infection. *Sex Transm Infect.* 1999;75(6):398-402.
44. Green J, Kocsis A. Psychological factors in recurrent genital herpes. *Genitourin Med.* 1997;73(4):253-8.
45. Nilsen AE, Aasen T, Halsos AM, Kinge BR, Tjøtta EA, Wikstrom K, et al. Efficacy of oral acyclovir in the treatment of initial and recurrent genital herpes. *Lancet.* 1982;2(8298):571-3.
46. Spruance SL, Tyring SK, DeGregorio B, Miller C, Beutner K. A large-scale, placebo-controlled, dose-ranging trial of peroral valaciclovir for episodic treatment of recurrent herpes genitalis. Valaciclovir HSV Study Group. *Arch Intern Med.* 1996;156(15):1729-35.
47. Sacks SL, Aoki FY, Diaz-Mitoma F, Sellors J, Shafran SD. Patient-initiated, twice-daily oral famciclovir for early recurrent genital herpes. A randomized, double-blind multicenter trial. Canadian Famciclovir Study Group. *JAMA.* 1996;276(1):44-9.
48. Spruance SL, Overall JC, Jr., Kern ER, Krueger GG, Pliam V, Miller W. The natural history of recurrent herpes simplex labialis: implications for antiviral therapy. *N Engl J Med.* 1977;297(2):69-75.
49. Wald A, Carrell D, Remington M, Kexel E, Zeh J, Corey L. Two-day regimen of acyclovir for treatment of recurrent genital herpes simplex virus type 2 infection. *Clin Infect Dis.* 2002;34(7):944-8.
50. Bodsworth N, Bloch M, McNulty A, Denham I, Doong N, Trottier S, et al. 2-day versus 5-day famciclovir as treatment of recurrences of genital herpes: results of the FaST study. *Sex Health.* 2008;5(3):219-25.
51. Aoki FY, Tyring S, Diaz-Mitoma F, Gross G, Gao J, Hamed K. Single-day, patient-initiated famciclovir therapy for recurrent genital herpes: a randomized, double-blind, placebo-controlled trial. *Clin Infect Dis.* 2006;42(1):8-13.
52. Abudalu M, Tyring S, Koltun W, Bodsworth N, Hamed K. Single-day, patient-initiated famciclovir therapy versus 3-day valacyclovir regimen for recurrent genital herpes: a randomized, double-blind, comparative trial. *Clin Infect Dis.* 2008;47(5):651-8.
53. Leone PA, Trottier S, Miller JM. Valacyclovir for episodic treatment of genital herpes: a shorter 3-day treatment course compared with 5-day treatment. *Clin Infect Dis.* 2002;34(7):958-62.
54. Strand A, Patel R, Wulf HC, Coates KM. Aborted genital herpes simplex virus lesions: findings from a randomised controlled trial with valaciclovir. *Sex Transm Infect.* 2002;78(6):435-9.
55. Mertz GJ, Jones CC, Mills J, Fife KH, Lemon SM, Stapleton JT, et al. Long-term acyclovir suppression of frequently recurring genital herpes simplex virus infection. A multicenter double-blind trial. *JAMA.* 1988;260(2):201-6.
56. Mindel A, Faherty A, Carney O, Patou G, Freris M, Williams P. Dosage and safety of long-term suppressive acyclovir therapy for recurrent genital herpes. *Lancet.* 1988;1(8591):926-8.

57. Lebrun-Vignes B, Bouzamondo A, Dupuy A, Guillaume JC, Lechat P, Chosidow O. A meta-analysis to assess the efficacy of oral antiviral treatment to prevent genital herpes outbreaks. *J Am Acad Dermatol.* 2007;57(2):238-46.
58. Wasserheit JN. Epidemiological synergy. Interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis.* 1992;19(2):61-77.
59. Hook EW, 3rd, Cannon RO, Nahmias AJ, Lee FF, Campbell CH, Jr., Glasser D, et al. Herpes simplex virus infection as a risk factor for human immunodeficiency virus infection in heterosexuals. *J Infect Dis.* 1992;165(2):251-5.
60. Mosca JD, Bednarik DP, Raj NB, Rosen CA, Sodroski JG, Haseltine WA, et al. Activation of human immunodeficiency virus by herpesvirus infection: identification of a region within the long terminal repeat that responds to a trans-acting factor encoded by herpes simplex virus 1. *Proceedings of the National Academy of Sciences of the United States of America.* 1987;84(21):7408-12.
61. Ostrove JM, Leonard J, Weck KE, Rabson AB, Gendelman HE. Activation of the human immunodeficiency virus by herpes simplex virus type 1. *Journal of virology.* 1987;61(12):3726-32.
62. Golden MP, Kim S, Hammer SM, Ladd EA, Schaffer PA, DeLuca N, et al. Activation of human immunodeficiency virus by herpes simplex virus. *J Infect Dis.* 1992;166(3):494-9.
63. Margolis DM, Rabson AB, Straus SE, Ostrove JM. Transactivation of the HIV-1 LTR by HSV-1 immediate-early genes. *Virology.* 1992;186(2):788-91.
64. Koelle DM, Abbo H, Peck A, Ziegweid K, Corey L. Direct recovery of herpes simplex virus (HSV)-specific T lymphocyte clones from recurrent genital HSV-2 lesions. *J Infect Dis.* 1994;169(5):956-61.
65. Kreiss J, Carael M, Meheus A. Role of sexually transmitted diseases in transmitting human immunodeficiency virus. *Genitourin Med.* 1988;64(1):1-2.
66. Kreiss JK, Coombs R, Plummer F, Holmes KK, Nikora B, Cameron W, et al. Isolation of human immunodeficiency virus from genital ulcers in Nairobi prostitutes. *J Infect Dis.* 1989;160(3):380-4.
67. Schacker T, Ryncarz AJ, Goddard J, Diem K, Shaughnessy M, Corey L. Frequent recovery of HIV-1 from genital herpes simplex virus lesions in HIV-1-infected men. *Jama.* 1998;280(1):61-6.
68. Nagot N, Ouedraogo A, Foulongne V, Konate I, Weiss HA, Vergne L, et al. Reduction of HIV-1 RNA levels with therapy to suppress herpes simplex virus. *N Engl J Med.* 2007;356(8):790-9.
69. Todd J, Grosskurth H, Changalucha J, Obasi A, Moshafiq F, Balira R, et al. Risk factors influencing HIV infection incidence in a rural African population: a nested case-control study. *J Infect Dis.* 2006;193(3):458-66.
70. Freeman EE, Weiss HA, Glynn JR, Cross PL, Whitworth JA, Hayes RJ. Herpes simplex virus 2 infection increases HIV acquisition in men and women: systematic review and meta-analysis of longitudinal studies. *Aids.* 2006;20(1):73-83.
71. Chen CY, Ballard RC, Beck-Sague CM, Dangor Y, Radebe F, Schmid S, et al. Human immunodeficiency virus infection and genital ulcer disease in South Africa: the herpetic connection. *Sex Transm Dis.* 2000;27(1):21-9.
72. Bagdades EK, Pillay D, Squire SB, O'Neil C, Johnson MA, Griffiths PD. Relationship between herpes simplex virus ulceration and CD4+ cell counts in patients with HIV infection. *Aids.* 1992;6(11):1317-20.
73. Wright PW, Hoesley CJ, Squires KE, Croom-Rivers A, Weiss HL, Gnann JW, Jr. A prospective study of genital herpes simplex virus type 2 infection in human immunodeficiency virus type 1 (HIV-1)-seropositive women: correlations with CD4 cell count and plasma HIV-1 RNA level. *Clin Infect Dis.* 2003;36(2):207-11.
74. Quinnan GV, Jr., Masur H, Rook AH, Armstrong G, Frederick WR, Epstein J, et al. Herpesvirus infections in the acquired immune deficiency syndrome. *Jama.* 1984;252(1):72-7.
75. Maier JA, Bergman A, Ross MG. Acquired immunodeficiency syndrome manifested by chronic primary genital herpes. *Am J Obstet Gynecol.* 1986;155(4):756-8.

76. Leung DT, Sacks SL. Current recommendations for the treatment of genital herpes. *Drugs*. 2000;60(6):1329-52.
77. Drew WL BW, Erlich KS et al. Chapter 34: Management of herpes virus infection (cytomegalovirus, herpes simplex virus and varicella zoster virus). In: Volberding P, Greene W, Joep L et al. *The medical management of AIDS*. 9th ed. London: Saunders, 2012, p.433-454.
78. Aoki FY. Management of genital herpes in HIV-infected patients. *Herpes*. 2001;8(2):41-5.
79. Foley E, Patel R. Treatment of genital herpes infections in HIV-infected patients. *Journal of HIV therapy*. 2004;9(1):14-8.
80. Marks GL, Nolan PE, Erlich KS, Ellis MN. Mucocutaneous dissemination of acyclovir-resistant herpes simplex virus in a patient with AIDS. *Reviews of infectious diseases*. 1989;11(3):474-6.
81. Celum C, Wald A, Hughes J, Sanchez J, Reid S, Delany-Moretlwe S, et al. Effect of aciclovir on HIV-1 acquisition in herpes simplex virus 2 seropositive women and men who have sex with men: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2008;371(9630):2109-19.
82. Celum C, Wald A, Lingappa JR, Magaret AS, Wang RS, Mugo N, et al. Acyclovir and transmission of HIV-1 from persons infected with HIV-1 and HSV-2. *N Engl J Med*. 2010;362(5):427-39.
83. Sacks SL, Wanklin RJ, Reece DE, Hicks KA, Tyler KL, Coen DM. Progressive esophagitis from acyclovir-resistant herpes simplex. Clinical roles for DNA polymerase mutants and viral heterogeneity? *Ann Intern Med*. 1989;111(11):893-9.
84. Gateley A, Gander RM, Johnson PC, Kit S, Otsuka H, Kohl S. Herpes simplex virus type 2 meningoencephalitis resistant to acyclovir in a patient with AIDS. *J Infect Dis*. 1990;161(4):711-5.
85. Englund JA, Zimmerman ME, Swierkosz EM, Goodman JL, Scholl DR, Balfour HH, Jr. Herpes simplex virus resistant to acyclovir. A study in a tertiary care center. *Ann Intern Med*. 1990;112(6):416-22.
86. Ratnam I, Chiu C, Kandala NB, Easterbrook PJ. Incidence and risk factors for immune reconstitution inflammatory syndrome in an ethnically diverse HIV type 1-infected cohort. *Clin Infect Dis*. 2006;42(3):418-27.
87. Fox PA, Barton SE, Francis N, Youle M, Henderson DC, Pillay D, et al. Chronic erosive herpes simplex virus infection of the penis, a possible immune reconstitution disease. *HIV medicine*. 1999;1(1):10-8.
88. Lalezari J, Schacker T, Feinberg J, Gathe J, Lee S, Cheung T, et al. A randomized, double-blind, placebo-controlled trial of cidofovir gel for the treatment of acyclovir-unresponsive mucocutaneous herpes simplex virus infection in patients with AIDS. *J Infect Dis*. 1997;176(4):892-8.
89. CDC - 2015 Sexually Transmitted Diseases Treatment Guidelines - Genital HSV Infections. <http://www.cdc.gov/std/tg2015/herpes.htm> (accessed 2nd September 2016).
90. DeJesus E, Wald A, Warren T, Schacker TW, Trottier S, Shahmanesh M, et al. Valacyclovir for the suppression of recurrent genital herpes in human immunodeficiency virus-infected subjects. *J Infect Dis*. 2003;188(7):1009-16.
91. Warren T, Harris J, Brennan CA. Efficacy and safety of valacyclovir for the suppression and episodic treatment of herpes simplex virus in patients with HIV. *Clin Infect Dis*. 2004;39 Suppl 5:S258-66.
92. Delany S, Mlaba N, Clayton T, Akpomiemie G, Capovilla A, Legoff J, et al. Impact of aciclovir on genital and plasma HIV-1 RNA in HSV-2/HIV-1 co-infected women: a randomized placebo-controlled trial in South Africa. *AIDS*. 2009;23(4):461-9.
93. Stranska R, Schuurman R, Nienhuis E, Goedegebuure IW, Polman M, Weel JF, et al. Survey of acyclovir-resistant herpes simplex virus in the Netherlands: prevalence and characterization. *J Clin Virol*. 2005;32(1):7-18.
94. Hill EL, Hunter GA, Ellis MN. In vitro and in vivo characterization of herpes simplex virus clinical isolates recovered from patients infected with human immunodeficiency virus. *Antimicrobial agents and chemotherapy*. 1991;35(11):2322-8.

95. Safrin S. Treatment of acyclovir-resistant herpes simplex virus infections in patients with AIDS. *Journal of acquired immune deficiency syndromes*. 1992;5 Suppl 1:S29-32.
96. Gnann J, Davis M, Harden E, et al. Resistance T force on H. acyclovir-resistant HSV from HIV-infected individuals. Population surveillance and in vitro characterization of isolates (abstract). New Orleans, USA, 38th Annual International Disease Society of America (IDSA) Meeting, 2000.
97. Engel JP, Englund JA, Fletcher CV, Hill EL. Treatment of resistant herpes simplex virus with continuous-infusion acyclovir. *Jama*. 1990;263(12):1662-4.
98. Sheffield JS, Hollier LM, Hill JB, Stuart GS, Wendel GD. Acyclovir prophylaxis to prevent herpes simplex virus recurrence at delivery: a systematic review. *Obstet Gynecol*. 2003;102(6):1396-403.
99. Watts DH, Brown ZA, Money D, Selke S, Huang ML, Sacks SL, et al. A double-blind, randomized, placebo-controlled trial of acyclovir in late pregnancy for the reduction of herpes simplex virus shedding and cesarean delivery. *Am J Obstet Gynecol*. 2003;188(3):836-43.
100. Scott LL, Hollier LM, McIntire D, Sanchez PJ, Jackson GL, Wendel GD, Jr. Acyclovir suppression to prevent recurrent genital herpes at delivery. *Infect Dis Obstet Gynecol*. 2002;10(2):71-7.
101. Brocklehurst P, Kinghorn G, Carney O, Helsen K, Ross E, Ellis E, et al. A randomised placebo controlled trial of suppressive acyclovir in late pregnancy in women with recurrent genital herpes infection. *Br J Obstet Gynaecol*. 1998;105(3):275-80.
102. Scott LL, Sanchez PJ, Jackson GL, Zeray F, Wendel GD, Jr. Acyclovir suppression to prevent cesarean delivery after first-episode genital herpes. *Obstet Gynecol*. 1996;87(1):69-73.
103. Braig S, Luton D, Sibony O, Edlinger C, Boissinot C, Blot P, et al. Acyclovir prophylaxis in late pregnancy prevents recurrent genital herpes and viral shedding. *Eur J Obstet Gynecol Reprod Biol*. 2001;96(1):55-8.
104. Hollier LM, Wendel GD. Third trimester antiviral prophylaxis for preventing maternal genital herpes simplex virus (HSV) recurrences and neonatal infection. *Cochrane Database Syst Rev*. 2008(1):CD004946.
105. Gardella C, Brown ZA, Wald A, Morrow RA, Selke S, Krantz E, et al. Poor correlation between genital lesions and detection of herpes simplex virus in women in labor. *Obstet Gynecol*. 2005;106(2):268-74.
106. Chen KT, Segu M, Lumey LH, Kuhn L, Carter RJ, Bulterys M, et al. Genital herpes simplex virus infection and perinatal transmission of human immunodeficiency virus. *Obstet Gynecol*. 2005;106(6):1341-8.
107. Poeran J, Wildschut H, Gaytant M, Galama J, Steegers E, van der Meijden W. The incidence of neonatal herpes in The Netherlands. *J Clin Virol*. 2008;42(4):321-5.
108. Acyclovir and Valacyclovir in Pregnancy Registry final report. April 1999. <http://pregnancyregistry.gsk.com/acyclovir.html> (accessed 2nd September 2016).
109. Andrews WW, Kimberlin DF, Whitley R, Cliver S, Ramsey PS, Deeter R. Valacyclovir therapy to reduce recurrent genital herpes in pregnant women. *Am J Obstet Gynecol*. 2006;194(3):774-81.
110. Sheffield JS, Hill JB, Hollier LM, Laibl VR, Roberts SW, Sanchez PJ, et al. Valacyclovir prophylaxis to prevent recurrent herpes at delivery: a randomized clinical trial. *Obstet Gynecol*. 2006;108(1):141-7.

Appendices:

Search Strategy:

A literature search was performed using PubMed/MEDLINE, Google, The Cochrane Library and relevant guidelines from January 1981 to October 2016. A MEDLINE/PubMed search was carried out from January 1981 to October 2016 using the following search terms/Medical Subject Headings (MeSH): “HSV OR herpes”, “genital ulcer OR genital ulcers” and “pregnancy complications: infectious”. The search was limited to humans and the English language. For some specific recommendations, an additional MEDLINE/PubMed search was performed when necessary. A Google search was performed in October 2016 with the search term “HSV guideline OR HSV guidelines” and all relevant documents of the first 150 search results were reviewed. A search of The Cochrane Library included the Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effects and Cochrane Central Register of Controlled Trials. The following guidelines were reviewed in detail: 2010 European guideline for the management of genital herpes; 2014 BASHH guidance on the management of genital herpes; USA Centers for Disease Control and Prevention 2015 STD Treatment Guidelines; 2014 BASHH/RCOG Joint Guideline on the management of genital herpes in pregnancy; and the 2016 WHO guidelines on the treatment of genital herpes simplex viruses. The members of the guideline development group selected studies relevant to the scope of the guideline. Article titles and abstracts were reviewed and if relevant the full-text article obtained. Priority was given to randomised controlled trial and systematic review evidence and, where possible, recommendations were made and graded on the basis of the best available evidence. In areas where evidence is lacking, recommendations based on consensus opinion within the writing group have been made.

Grading of Evidence:

For details of the Tables of levels of evidence and grading of recommendations please see:

http://www.iusti.org/regions/Europe/pdf/2013/Levels_of_Evidence.pdf

Statement on declarations of interest:

None to declare